

Use Foster[®] First Defense[®] in: Medical Related facilities, Nursing homes, Day care centers, Food processing plants, Restaurants, USDA inspected food processing facilities, Athletic facilities, Correctional facilities, Factories, Hotels, Schools, Airplanes, Buses, Emergency vehicles, Ships, Taxis, Trailers, Trains, Transportation terminals.

Foster[®] First Defense[®] delivers non-acid disinfectant and cleaning performance. Use on washable hard, non-porous surfaces of tables, chairs, countertops, floors, walls, coils and drain pans of air conditioning and refrigeration equipment and heat pumps, glazed tiles, ultrasonic baths, hard, non-porous athletic mats, wrestling mats, gymnastic mats, exercise equipment and training tables, non-critical medical equipment surfaces: examining tables, lamps, resuscitators, stretchers, stethoscopes, wheel chairs and other hard, non-porous surfaces made of: metal, plastic, stainless steel, finished woodwork and upholstery, vinyl and plastic. Use this product to clean and disinfect hard, non-porous personal protective safety equipment, protective headgear, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, hearing protectors, and ear muffs. Rinse all equipment that comes in prolonged contact with skin with warm water and allow to air dry before reuse. For use on soft surfaces such as upholstery, fabric, furniture, gym bags, and back packs.

Foster[®] First Defense[®] is a one-step² Hospital use disinfectant⁵, Bactericidal tested according to the current AOAC Use-Dilution Test Method, Fungicidal according to the AOAC Fungicidal Test and Virucidal* tested according to the virucidal qualification modified in the presence of 5% organic serum against:

BACTERIA - Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Acinetobacter baumannii, Burkholderia cepacia, Campylobacter jejuni, Corynebacterium ammoniagenes, Enterobacter aerogenes, Enterobacter cloacae NDM-1, Enterococcus faecalis, Enterococcus faecalis - Vancomycin Resistant (VRE), Escherichia coli (E. coli), Escherichia coli, New Delhi Metallo-Beta Lactamase (NDM-1), ESBL Escherichia coli, Klebsiella pneumoniae, Klebsiella pneumoniae - NDM-1 positive, Legionella pneumophila, Listeria monocytogenes (Listeria), Salmonella schottmuelieri (Salmonella), Salmonella typhi (Salmonella), Serratia marcescens, Shigella dysenteriae, Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRSA) (NRS384) (USA300), Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRSA) (NRS123) (USA400), Staphylococcus aureus - Methicillin-Resistant (MRSA), Staphylococcus aureus Multi-Drug Resistant, Staphylococcus aureus - Vancomycin Intermediate Resistant - (ISA), Streptococcus pyogenes (Strep), Vibrio cholerae.

FUNGI - Aspergillus niger, Aspergillus fumigatus, Trichophyton interdigitale.

VIROUSES - *Hepatitis B Virus (HBV), *Hepatitis C Virus (HCV), *Herpes Simplex Virus Type 1, *Herpes Simplex Virus Type 2, *HIV-1 (AIDS Virus), *Human Coronavirus, *Influenza Virus Type A / H1N1, *Influenza Virus Type A / H3N2 (Influenza), *2013 Influenza A Virus H7N9, *Respiratory Syncytial Virus (RSV), *SARS Associated Coronavirus (SARS), *SARS-Related Coronavirus 2 (SARS-CoV-2), cause of COVID-19, *Vaccinia, *Norwalk Virus (Norovirus), *Rotavirus.

ANIMAL VIROUSES - Avian Infectious Bronchitis Virus, Avian Influenza Virus H3N2, Avian Influenza Virus H5N1, Canine Distemper Virus, Feline Calicivirus, Murine Norovirus², Newcastle's Disease Virus, Pseudorabies Virus.

Foster[®] First Defense[®] is a proven soft surface sanitizer^{11,12} against: Staphylococcus aureus, Enterobacter aerogenes.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. You may also contact the poison control center at 1-800-222-1222 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Disinfectant

Ready-to-Use

Cleaner • Fungicide • Mildewstat • Virucide* • Deodorizer
For Homes, Restaurants, Hospitals, Institutional and Industrial Use

DISINFECTS 99.9% SARS-COV-2² Virus the cause of COVID-19

Effective against SARS-Related Coronavirus 2² (SARS-COV-2) IN 1 MINUTE

Kills Cold and Flu Virus*^{2,13}

- One-Step Disinfectant^{2,5} cleaner that is effective against a broad spectrum of bacteria, is virucidal*, fungicidal, and inhibits the growth of mold and mildew, and their odors, when used as directed.
- Use to remove odors caused by flooding and smoke from fires.
- Kills 99.9% of germs and bacteria, including the Flu Virus¹³, while cleaning to a virtually streak free finish.
- Proven effective soft surface sanitizer^{11,12}.

Active Ingredients:

Octyl decyl dimethyl ammonium chloride0.025%
Dioctyl dimethyl ammonium chloride.....0.010%
Didecyl dimethyl ammonium chloride.....0.015%
Alkyl (C₁₄, 50%; C₁₂, 40%; C₁₆, 10%)
dimethyl benzyl ammonium chloride.....0.034%
Other Ingredients.....99.916%
TOTAL.....100.000%

**KEEP OUT OF REACH
OF CHILDREN**

**FOR EMERGENCY MEDICAL
INFORMATION: 1-888-853-1758**

CAUTION

**See Left Panel for Precautionary
Statements & First Aid.**



SOLD BY:
H.B. Fuller Construction Products Inc.
1105 South Frontenac Street
Aurora, IL 60504-6451, U.S.A.
PH 800.832.9002 | fosterproducts.com

40010351 Rev Z R0523

Foster, First Defense and the Foster design are trademarks of H.B. Fuller.

MANUFACTURED in USA

NET CONTENTS:

5 Gallons (41 lbs.) / 18.92 L (18.59 KG)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Before use in a federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD AND MILDEW CONTROL
DIRECTIONS: Apply Foster[®] First Defense[®] undiluted to hard, non-porous surfaces, thoroughly wetting the surfaces with a cloth, mop, sponge, sprayer or by immersion. Treated surfaces must remain visibly wet for 10 minutes. **For Influenza Virus Type A, Human Coronavirus and SARS-CoV-2**, treated surfaces must remain visibly wet for 1 minute. Wipe dry with a cloth, sponge or mop or allow to air dry. For visibly soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface. Avoid breathing spray. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes. **For fungicidal activity:** Foster[®] First Defense[®] is an effective one-step fungicide against Trichophyton interdigitale when used on hard nonporous surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. Allow surface to remain visibly wet for 10 minutes. **For mold and mildew:** Repeat treatment every 7 days or more often if new growth appears.

This product is not to be used as a terminal sterilant / high level disinfectant on any surface or instrument that is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

***KILLS HIV-1 AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECT PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings with which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: Personal Protection:** Clean up must always be done wearing protective gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of Foster[®] First Defense[®]. **Contact Time:** Leave surfaces visibly wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV. Use a 10 minute contact time for disinfection against all other viruses, bacteria and fungi claimed. **Disposal of Infectious Material:** Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

ULTRASONIC BATH DISINFECTANT / VIRUCIDAL* DIRECTIONS: Use Foster[®] First Defense[®] to disinfect hard nonporous non-critical objects compatible with Ultrasonic cleaning units. Pour Foster[®] First Defense[®] directly into bath chamber. Pre-clean visibly soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to the manufacturer's directions. Remove objects and wash with sterile water or allow to air dry. Use fresh product for each use.

FOR USE ON SEALED FINISHED FLOORS: Apply with a damp mop or autoscrubber. Allow surface to remain visibly wet for 10 minutes. Wipe or allow to air dry.

ANIMAL CARE / KENNELS DISINFECTION / VIRUCIDAL* DIRECTIONS: For cleaning and disinfecting hard nonporous surfaces: equipment used for feeding or watering animals, utensils, instruments, cages, kennels, stables, catteries, pens, stalls and etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, pens, stalls and etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water.

Saturate surfaces with Foster[®] First Defense[®] for a period of 10 minutes. Wipe or allow to air dry. Immerse all hard, non-porous equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Thoroughly scrub all treated surfaces, then rinse all surfaces that come in contact with food, including equipment used for feeding or watering, with potable water before reuse. Ventilate buildings, animal enclosures, vehicles and other closed spaces. Do not hose animals or employ equipment until treatment has been absorbed, set or dried.

TO CLEAN AND DEODORIZE: Apply undiluted to hard, non-porous, non-food surfaces with a cloth, mop, sponge or sprayer. Wipe or allow to air dry. Rinse all surfaces that come in contact with food such as countertops, exteriors of appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

TO SANITIZE SOFT SURFACES¹¹: Hold bottle 6" - 8" from surface and spray until fabric is wet. Fabric must remain visibly wet for 5 minutes. Allow fabric to air dry. Visibly soiled areas must be cleaned prior to sanitizing.

WATER DAMAGE RESTORATION AGAINST ODOR-CAUSING BACTERIA AND FUNGI FOR HOME, INSTITUTIONAL AND INDUSTRIAL USE: (Not for use in California).

This product is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, subfloors, drywall, trim, and frame lumber, tackless strip and paneling. Saturate the affected materials with enough product to remain visibly wet for at least 10 minutes. **Sewer Back up and River Flooding:** During mitigation procedures remove visible filth or heavy soil along with non-salvageable materials. Saturate all affected areas with a coarse spray device before and after cleaning and extraction. **Carpets, Carpet Cushion and Other Porous Materials such as Subfloors, Drywall, Trim and Frame Lumber, Tackless Strip and Paneling:** For water damage from a clean source, extract excess water. Test hidden area for colorfastness. Remove visible filth or heavy soil. Saturate all affected areas with a coarse spray device. Roll, brush or agitate into the materials and allow the materials to remain visibly wet for 10 minutes. Follow with a thorough extraction. Dry rapidly and thoroughly.

Special Instructions for Cleaning Carpet Against Odor Causing Bacteria: Use this product in industrial and institutional areas such as motels, hotels, and schools. Use on wet cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for colorfastness. Do not mix this product with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using this product, set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the manufacturer's directions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Open dumping is prohibited. Store in original container in areas inaccessible to children. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOOTNOTES

² On hard, non-porous, non-food surfaces

³ Tested in deionized water

⁵ When used according to the disinfection directions

¹¹ Spot treatment of soft surfaces.

¹² When used according to the sanitization for soft surfaces directions.

¹³ Influenza Virus Type A / H1N1, Influenza Virus Type A / H3N2 and Influenza A Virus H7N9

EPA Est. No.
63836-TX-001